

# **Roots of Alexander polynomials of random positive 3-braids**

Nathan Dunfield (University of Illinois)


joint with Giulio Tiozzo

Based on: arXiv:2402.06771


Slides at: <https://dunfield.info/slides/banff2024.pdf>

3-strand braid group:  $\text{Br}_3 = \langle \sigma_1, \sigma_2 \mid \sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2 \rangle$

$w = \sigma_1\sigma_2\sigma_1^{-2}\sigma_2\sigma_1^2\sigma_2^{-1}$ :




Braid closure:  $\hat{w}$



Alexander polynomial in  $\mathbb{Z}[t^{\pm 1}]$

$$\Delta_{\hat{w}}(t) = t^4 - 2t^3 + 3t^2 - 2t + 1$$



Dehornoy (2015): roots of  $\Delta_{\widehat{w}}$  for positive  $w \in \text{Br}_3$  are highly structured.




$$\deg \Delta_{\widehat{w}} = \#w - 2 \approx 760$$

69.3% of roots on  $S^1$

Comparison to roots of  $\Delta_{\widehat{w}}$  for a random braid in  $\{\sigma_1, \sigma_1^{-1}, \sigma_2, \sigma_2^{-1}\}$ .





Roots of  $\Delta_{\hat{w}}$  for 2,500 positive braids with mean  $\#w \approx 500$  and std. dev. 170.



$w$  chosen randomly  
with  $\sigma_1$  and  $\sigma_2$  having  
probability  $1/2$

1.2 million roots shown

Distribution of roots of  $\Delta_{\hat{w}}$  on the top half of the circle.




For  $w \in \text{Br}_3$  let  $\nu_w$  be the prob. measure on  $\mathbb{C}$  unif. supported on the roots of  $\Delta_{\hat{w}}$ .

Generate a random walk  $w_n := g_1 g_2 \cdots g_n$  by picking  $(g_i)_{i \in \mathbb{N}} \in \{\sigma_1, \sigma_2\}^{\mathbb{N}}$  with respect to the uniform measure.

For  $w \in \text{Br}_3$  let  $\nu_w$  be the prob. measure on  $\mathbb{C}$  unif. supported on the roots of  $\Delta_{\hat{w}}$ .

Generate a random walk  $w_n := g_1 g_2 \cdots g_n$  by picking  $(g_i)_{i \in \mathbb{N}} \in \{\sigma_1, \sigma_2\}^{\mathbb{N}}$  with respect to the uniform measure.

**Conj.** There exists a compactly supported measure  $\nu_\infty$  on  $\mathbb{C}$  such that for a.e.  $w_n$  one has  $\nu_{w_n} \rightarrow \nu_\infty$  weakly. Moreover,  $\nu_\infty$  has the following properties:





Bureau rep  $B_t: \text{Br}_3 \rightarrow \text{GL}_2\mathbb{Z}[t^{\pm 1}]$  defined by  $\sigma_1 \mapsto \begin{pmatrix} -t & 1 \\ 0 & 1 \end{pmatrix}$  and  $\sigma_2 \mapsto \begin{pmatrix} 1 & 0 \\ t & -t \end{pmatrix}$

$$\Delta_{\hat{w}}(t) = \det(B_t(w) - 1) / (t^2 + t + 1)$$

For  $\bar{\mathbb{D}} = \{ |z| \leq 1 \}$ , take  $\rho_w: \bar{\mathbb{D}} \rightarrow \mathbb{R}_{\geq 0}$  to be the max abs. val. of an eig. val. of  $B_t(w)$ .



Key:  $\rho_w = 1$  at any root of  $\Delta_{\hat{w}}$  in  $\bar{\mathbb{D}}$ .

Burau rep  $B_t: \text{Br}_3 \rightarrow \text{GL}_2\mathbb{Z}[t^{\pm 1}]$  defined by  $\sigma_1 \mapsto \begin{pmatrix} -t & 1 \\ 0 & 1 \end{pmatrix}$  and  $\sigma_2 \mapsto \begin{pmatrix} 1 & 0 \\ t & -t \end{pmatrix}$

$$\Delta_{\hat{w}}(t) = \det(B_t(w) - 1) / (t^2 + t + 1)$$


For  $\bar{\mathbb{D}} = \{ |z| \leq 1 \}$ , take  $\rho_w: \bar{\mathbb{D}} \rightarrow \mathbb{R}_{\geq 0}$  to be the max abs. val. of an eig. val. of  $B_t(w)$ .

Key:  $\rho_w = 1$  at any root of  $\Delta_{\hat{w}}$  in  $\bar{\mathbb{D}}$ . Set  $R_w = \{ z \in \bar{\mathbb{D}} \mid \rho_w(z) = 1 \}$ .




**Thm.** For any positive  $w \neq \sigma_i^k$ , set  $R_w$  contains the arc  $A_R := \{t = e^{i\theta} \mid |\theta| < 2\pi/3\}$ , is disjoint from the set  $\mathcal{T}$ , and meets  $(-1, 1)$  in a single point.

**Thm.** For any positive  $w \neq \sigma_i^k$ , at least  $\frac{2}{3}(\deg(\Delta_{\hat{w}}) - 1)$  of the roots of  $\Delta_{\hat{w}}$  occur on this arc.



**Thm.** For any positive  $w \neq \sigma_i^k$ , set  $R_w$  contains the arc  $A_R := \{t = e^{i\theta} \mid |\theta| < 2\pi/3\}$ , is disjoint from the set  $\mathcal{T}$ , and meets  $(-1, 1)$  in a single point.

**Thm.** For any positive  $w \neq \sigma_i^k$ , at least  $\frac{2}{3}(\deg(\Delta_{\hat{w}}) - 1)$  of the roots of  $\Delta_{\hat{w}}$  occur on this arc.



**Thm.** Set  $A_L := \{t = e^{i\theta} \mid |\theta - \pi| < \pi/3\}$ . For a.e. random walk  $w_n$ , asymptotically the portion of roots on  $A_L$  is  $\geq (7 - 3\sqrt{5})/12 \approx 2.4\%$ .

**Thm.** The signature  $\sigma_{\hat{w}}(-1)$  obeys a central limit theorem with positive drift  $(5 - \sqrt{5})/4$ .

Lyapunov exponent  $\lambda(t) := \lim_{n \rightarrow \infty} \frac{1}{n} \int_{\text{Br}_3} \log \|B_t(g)\| d\mu^{*n}(g)$  where  $\mu$  is the uniform measure on  $\{\sigma_1, \sigma_2\}$ .


$$\chi(t) := \max \{ \lambda(t), \log |t|, 0 \}.$$

Bifurcation measure:  $\nu_{bif} := \Delta \chi$

**Conj.**  $\nu_{W_n} \rightarrow \nu_{bif}$

Motivated by Deroin-Dujardin.

We have some partial results towards this conjecture.





## Open questions:

Prove the conjecture!

$n$ -strand braids

Non-positive braids?



*Ribbon concordances and slice obstructions: experiments and examples*

Forthcoming work with Sherry Gong

- ▶ 352 million knots with  $\leq 19$  crossings [Burton]
- ▶ 1.6 million are slice
- ▶ 350.4 million are not slice
- ▶  $< 13,000$  unknown (0.004%)